Glucose Tolerance (2 Hour) w/o Urine

Order Name: GTT 2 HR
Test Number: 2016800
Revision Date: 05/20/2021

TEST NAME	METHODOLOGY	LOINC CODE
Glucose Fasting	Hexokinase	1558-6
Glucose 1 Hour Tolerance	Hexokinase	20438-8
Glucose 2 Hour Total	Hexokinase	20436-2

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5)	Plasma	Sodium fluoride (gray top)	Refrigerated
Alternate 1	1 mL (0.5)	Plasma	Lithium Heparin PST (Light Green Top)	Refrigerated
Alternate 2	1 mL (0.5)	Serum	Clot Activator SST	Refrigerated
Instructions	Fasting overnight and during test. Patient may have water. Draw fasting specimen. Adults: Give 75 grams of glucola. Children: Adjusted amount of glucola to be calculated by lab. Call (918) 744-2500. Stability: Ambient 8 hours, Refrigerated 7 days, Frozen 1 month Fluoride preserved plasma samples are stable at room temperature for 24 hours			

GENERAL INFORMATION		
Testing Schedule	Mon - Fri	
Expected TAT	1-2 days	
Clinical Use	Criteria for Diagnosis of Diabetes from the American Diabetes Association recommends Random plasma glucose >200 mg/dl with symptoms (polyuria, polydypsia, and unexplained weight loss) repeated to confirm on subsequent day, or Fasting plasma glucose>126 mg/dl repeated to confirm, or 2hr plasma glucose <200 mg/dl post 75g glucose challenge repeated to confirm.	
Notes	For more information on this test, access our "Specialized Tests" section.	
CPT Code(s)	82951	
Lab Section	Chemistry	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025