Labcorp Oklahoma, Inc. Test Directory

Norpace (Disopyramide)

Order Name: NORPACE
Test Number: 4003260
Revision Date: 12/12/2022

TEST NAME			METHODOLOGY	LOINC CODE
Norpace (Disopyramide)			Immunoassay (IA)	
SPECIMEN REQUIR	EMENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1mL (0.3 mL)	Serum	Clot Activator (Red Top, No-Gel)	Room Temperature
Alternate 1	1mL (0.3 mL)	Plasma	EDTA (Lavender Top)	Room Temperature
Instructions	Specimen Type: Red-top tube, lavender-top (EDTA) tube OR green-top (heparin) tube. DO NOT USE A GEL-BARRIER TUBE. The use of gelbarrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant. Specimen Storage: Room Temperature Specimen Collection: Transfer separated serum or plasma to a plastic transport tube. Specimen Stability: Ambient: 14 days, Refrigerated: 14 days, Frozen: 14 days			

GENERAL INFORMATION		
Expected TAT	2 - 3 days working day.	
Clinical Use	Disopyramide is useful in treating patients with cardiac arrhythmias and tachcardia. Therapeutic drug monitoring is useful to optimize dose and avoid toxicity.	
Performing Labcorp Test Code	007864	
Notes	Labcorp Test Code: 007864	
CPT Code(s)	80299	
Lab Section	Reference Lab	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025