

## Procainamide

Order Name: **PROCAINAMI**  
Test Number: 3621850  
Revision Date: 12/12/2022

TEST NAME	METHODOLOGY	LOINC CODE
Procainamide	Immunoassay (IA)	

### SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1mL (0.3 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated
Alternate 1	1mL (0.3 mL)	Plasma	Sodium Heparin (Green Top / No-Gel)	Refrigerated

**Instructions**

**Collect as a trough just prior to next dose**

**Notes:** 0.3 mL (Note: This volume Does NOT allow for repeat testing.)

**Specimen Type:** Red-top tube OR green-top (heparin) tube. **DO NOT USE A GEL-BARRIER TUBE.** The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Depending on the specimen volume and storage time, the decrease in drug level due to absorption may be clinically significant.

**Container Detail:** No Gel

**Specimen Storage:** Refrigerated

**Specimen Collection:** Transfer separated serum or plasma to a plastic transport tube. Oral treatment: peak: 75 minutes after dose; trough: immediately before next dose. I.V. treatment: immediately after loading dose; 2, 6, 12, and 24 hours after starting I.V. maintenance.

**Specimen Stability:** Ambient: 2 days, Refrigerated : 14 days, Frozen: 14 days

### GENERAL INFORMATION

Expected TAT	2-3 Days
Clinical Use	Procainamide is used to treat a variety of atrial and ventricular arrhythmias. Metabolism of procainamide produces an active metabolite N-Acetylprocainamide (NAPA). The concentration of both procainamide and NAPA levels are monitored to assure adequate therapeutic levels of procainamide are achieved and to avoid toxicity.
Performing Labcorp Test Code	007252
Notes	Labcorp Test Code: 007252
CPT Code(s)	80192
Lab Section	Reference Lab