Metapneumovirus PCR

Order Name: METAPNEUMO Test Number: 5575875 Revision Date: 10/01/2022

TEST NAME		ME	METHODOLOGY		
Metapneumovirus PCI	R	Po	Polymerase Chain Reaction		
SPECIMEN REQUIREMENTS					
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	3 mL (1 mL)	Swab	Mini-Flocked Swab in Universal Transport Media (UTM)	Refrigerated	
Alternate 1	3 mL (1 mL)	Nasal Wash	Sterile Screwtop Container	Refrigerated	
Alternate 2	3 mL (1 mL)	Bronchial lavage/wash	Sterile Screwtop Container	Refrigerated	
Instructions	(Comes as a kit: Supply# 5077 Keep swabs refrigerated, Stabi Also acceptable 3mL(1mL) BA	The preferred specimen is mini-Flocked Swab in Universal Transport Media (UTM) Comes as a kit: Supply# 50775), BD Viral Transport Media (VTM) or M4. Keep swabs refrigerated, Stability 7 days refrigerated. Also acceptable 3mL(1mL) BAL or NP/Nasal/Tracheal Aspirate Sterile Screwtop tube Refrigerated. Jnacceptable Spcimens: Eswabs, Caclium Alginate or Wooden Shaft Swabs.			
GENERAL INFORMATION					
Testing Schedule	In season: Daily. Out	In season: Daily. Out of season: Mon, Wed, Fri			
Expected TAT	1-3 Days	1-3 Days			
Notes	PCR detection of the	PCR detection of the pathogen's RNA or DNA will provide a more sensitive and specific method when compared to the DFA method.			

 CPT Code(s)
 87798

 Lab Section
 Molecular Diagnostic