## Labcorp Oklahoma, Inc. Test Directory

## Epi proColon

Order Name: Epi proColon

Test Number: 6904831 Revision Date: 12/12/2022

		METHODOLOGY	LOINC CODE
		Polymerase Chain Reaction	65810-4
EMENTS			
Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Two 3.5 mL	Plasma	EDTA (Lavender Top)	Frozen
Notes: One plastic transport tube (containing at least 3.5 mL) of frozen Plasma. Note: This volume Does NOT allow for repeat testing.  Specimen Type: 4-mL BD(R) purple-top (K2 EDTA) tube. Collect six tubes of whole blood in accordance with the collection procedure specified below.  Specimen Storage: FREEZE. The plasma sample could be stored frozen, at -15 (degrees)C to -25(degrees)C for up to 14 days.  Specimen Collection: Please label each final plasma collection tube with the word "Plasma". Testing will be delayed if the tube is not labeled properly. Blood Collection Blood should be collected according to your laboratory's procedure for venipuncture using ONLY a BD K2EDTA (purple top) 4 mL blood collection tube. Plasma preparation should be performed within four hours after blood is collected. Store, transport and ship plasma at frozen condition only. The plasma sample can be stored frozen at -15(degrees)C to -25(degrees)C for up to 14 days. NOTE: DO NOT FREEZE WHOLE BLOOD SAMPLES. Plasma Specimen Preparation Immediately following specimen collection, label all tubes with appropriate patient information (two patient identifiers, minimum). The brake setting of the Drucker 642e at LabCorp service center is at 3 or 4 (manufacturer setting). Centrifuge the six, BD K2EDTA (purple top) blood collection tubes for 10 minutes at 1600 plus or minus 90 rcf. [For conversion of RPM (revolutions per minute) to rcf (relative centrifugal force), refer to the centrifuge manufacturer's user manual.] Remove blood collection tubes from the centrifuge (Plasma sample will be rejected if it is hemolyzed. Patient will be contacted). Using a fresh six inch disposable transfer pipette, transfer plasma from three, 4 mL blood collection tubes to one 8.5 mL Sarstedt centrifuge tube (55.598.006). Repeat this process with the second set of three 4 mL blood tubes. Two 8.5 mL tubes will be collected from six, 4 mL blood collection tubes. Centrifuge plasma in the 8.5 mL centrifuge tube for 10 minutes at 1600 plus or minus 90 rcf. Using a fres			with the collection procedure specified  ses)C for up to 14 days.  be delayed if the tube is not labeled cture using ONLY a BD K2EDTA (purple collected. Store, transport and ship plasma up to 14 days. NOTE: DO NOT FREEZE ibel all tubes with appropriate patient center is at 3 or 4 (manufacturer setting).  Orcf. [For conversion of RPM (revolutions blood collection tubes from the centrifuge able transfer pipette, transfer plasma from ss with the second set of three 4 mL blood 3.5 mL centrifuge tube for 10 minutes at at 3.5 mL of plasma from one 8.5 mL 019). Repeat this process with the second 5(degrees)C. Stable for up to 14 days.
	Specimen Volume (min)  Two 3.5 mL  Notes: One plastic transport to Specimen Type: 4-mL BD(R) below.  Specimen Storage: FREEZE. Specimen Collection: Please properly. Blood Collection Blood top) 4 mL blood collection tube at frozen condition only. The p WHOLE BLOOD SAMPLES. Finformation (two patient identificant Centrifuge the six, BD K2EDT/per minute) to rcf (relative cent (Plasma sample will be rejected three, 4 mL blood collection tubes. Two 8.5 mL tubes will be 1600 plus or minus 90 rcf. Using centrifuge tube into a labeled 78.5 mL centrifuge tube. Two, 7 Note: Take care not to disturb	Two 3.5 mL  Plasma  Notes: One plastic transport tube (containing at least 3.5 m Specimen Type: 4-mL BD(R) purple-top (K2 EDTA) tube. Obelow.  Specimen Storage: FREEZE. The plasma sample could be Specimen Collection: Please label each final plasma colle properly. Blood Collection Blood should be collected accord top) 4 mL blood collection tube. Plasma preparation should at frozen condition only. The plasma sample can be stored WHOLE BLOOD SAMPLES. Plasma Specimen Preparation information (two patient identifiers, minimum). The brake se Centrifuge the six, BD K2EDTA (purple top) blood collection per minute) to rcf (relative centrifugal force), refer to the cer (Plasma sample will be rejected if it is hemolyzed. Patient we three, 4 mL blood collection tubes to one 8.5 mL Sarstedt of tubes. Two 8.5 mL tubes will be collected from six, 4 mL blood plus or minus 90 rcf. Using a fresh, six inch disposable centrifuge tube into a labeled 7.0 mL Sarstedt screw cap, fla 8.5 mL centrifuge tube. Two, 7.0 mL tubes will be collected. Note: Take care not to disturb or transfer the buffy coat (who	EMENTS  Specimen Volume (min) Specimen Type Specimen Container  Two 3.5 mL Plasma EDTA (Lavender Top)  Notes: One plastic transport tube (containing at least 3.5 mL) of frozen Plasma. Note: This volume Does I Specimen Type: 4-mL BD(R) purple-top (K2 EDTA) tube. Collect six tubes of whole blood in accordance below.  Specimen Storage: FREEZE. The plasma sample could be stored frozen, at -15 (degrees)C to -25 (degree Specimen Collection: Please label each final plasma collection tube with the word "Plasma". Testing will properly. Blood Collection Blood should be collected according to your laboratory's procedure for venipunc top) 4 mL blood collection tube. Plasma preparation should be performed within four hours after blood is c at frozen condition only. The plasma sample can be stored frozen at -15 (degrees)C to -25 (degrees)C for the UNIOLE BLOOD SAMPLES. Plasma Specimen Preparation Immediately following specimen collection, la information (two patient identifiers, minimum). The brake setting of the Drucker 642e at LabCorp service of Centrifuge the six, BD K2EDTA (purple top) blood collection tubes for 10 minutes at 1600 plus or minus 90 per minute) to rcf (relative centrifugal force), refer to the centrifuge manufacturer's user manual.] Remove (Plasma sample will be rejected if it is hemolyzed. Patient will be contacted). Using a fresh six inch dispose three, 4 mL blood collection tubes to one 8.5 mL Sarstedt centrifuge tube (55.598.006). Repeat this proces tubes. Two 8.5 mL tubes will be collected from six, 4 mL blood collection tubes. Centrifuge plasma in the 8 1600 plus or minus 90 rcf. Using a fresh, six inch disposable transfer pipette or serological pipette, transfer centrifuge tube into a labeled 7.0 mL Sarstedt screw cap, flat bottom purple frozen transport tube (62.550. 8.5 mL centrifuge tube. Two, 7.0 mL tubes will be collected. Ship and store plasma at -15 (degrees)C to -2

GENERAL INFORMATION		
Expected TAT	5 - 7 days	
Clinical Use	The Epi proColon test is indicated to screen adults of either sex, 50 years or older, defined as average risk for CRC, who have been offered and have a history of not completing CRC screening. Tests that are available and recommended in the USPSTF 2008 CRC screening guidelines should be offered and declined prior to offering the Epi proColon test. Patients with a positive Epi proColon test result should be referred for diagnostic colonoscopy. The Epi proColon test results should be used in combination with physician's assessment and individual risk factors in guiding patient management.	
Performing Labcorp Test Code	481160	
Notes	DNA is isolated from plasma specimen and treated with bisulfite. Real-time PCR is performed on ABI 7500 Fast Dx to detect methylated form of Septin 9 DNA. ?Labcorp Test Code: 481160	
CPT Code(s)	81327	
Lab Section	Reference Lab	

Specimen Stability: Ambient: Not Available, Refrigerated : Not Available, Frozen: Not Available

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025 Labcorp Oklahoma, Inc. Test Directory