## Labcorp Oklahoma, Inc. Test Directory

## **Anti-HMGCR Ab**

Order Name: HMGCR AB
Test Number: 5194380
Revision Date: 03/27/2025

		METHODOLOGY	LOINC CODE
		Chemiluminescence Assay	95726-6
REMENTS			
Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
2 mL (0.5 mL)	Serum	Clot Activator (SST or Red No-Gel)	Refrigerated
Specimen Type: gel-barrier to Specimen Storage: Refrigera Specimen Collection: Separa Special Instructions: Testing Specimen Stability: Ambients	ube or Red-top tube Ited ate serum from cells within one I It is performed at Esoterix Inc. Est 48 hours, Refrigerated: 14 day	nour of collection. Transfer to a plastic transport tub	e before shipping.
F	Specimen Volume (min)  2 mL (0.5 mL)  Notes: 1 mL (0.5 mL) Note: The Specimen Type: gel-barrier to Specimen Storage: Refrigeral Specimen Collection: Separate Special Instructions: Testing Specimen Stability: Ambients	Specimen Volume (min)  Specimen Type  2 mL (0.5 mL)  Serum  Notes: 1 mL (0.5 mL) Note: This volume Does not allow for re Specimen Type: gel-barrier tube or Red-top tube Specimen Storage: Refrigerated Specimen Collection: Separate serum from cells within one I Special Instructions: Testing is performed at Esoterix Inc. ES Specimen Stability: Ambient: 48 hours, Refrigerated: 14 day	Chemiluminescence Assay  REMENTS  Specimen Volume (min) Specimen Type Specimen Container  2 mL (0.5 mL) Serum Clot Activator (SST or Red No-Gel)  Notes: 1 mL (0.5 mL) Note: This volume Does not allow for repeat testing.)  Specimen Type: gel-barrier tube or Red-top tube

GENERAL INFORMATION		
Expected TAT	7 to 10 days	
Clinical Use	<ol> <li>Not all idiopathic inflammatory myopathy patients are positive for HMGCR antibodies.</li> <li>Results of this assay should be used in conjunction with clinical findings and other serological tests.</li> <li>Failure to adequately resuspend the HMGCR coated beads may yield lower values than if the beads are properly re-suspended.</li> <li>The performance characteristics of this assay have not been established for matrices other than serum.</li> <li>Bottai M, Tjarnlund A, Santoni G, et al. EULAR/ACR classification criteria for adult and juvenile idiopathic inflammatory myopathies and their major subgroups: a methodology report. RMD Open. 2017 Nov 14;3(2):e000507. 29177080</li> <li>Christopher-Stine L, Casciola-Rosen LA, Hong G, Chung T, Corse AM, Mammen AL. A novel autoantibody recognizing 200-kd and 100-kd proteins is associated with an immune-mediated necrotizing myopathy. Arthritis Rheum. 2010 Sep;62(9):2757-2766. 20496415</li> <li>Musset L, Allenbach Y, Benveniste O, et al. Anti-HMGCR antibodies as a biomarker for immune-mediated necrotizing myopathies: A history of statins and experience from a large international multi-center study. Autoimmun Rev. 2016 Oct;15(10):983-993. 27491568</li> </ol>	
Performing Labcorp Test Code	520385	
CPT Code(s)	83516	
Lab Section	Reference Lab	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025