Chlamydia/Gonococcus/Genital Mycoplasma Profile, NAA, Urine

Order Name: Ct Ng Myco UR

Test Number: 5195320 Revision Date: 09/12/2023

TEST NAME METHODOLOGY LOINC CODE

Chlamydia/Gonococcus/Genital Mycoplasma Profile, NAA, Urine

Nucleic acid amplification (NAA)

SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	30 mL (5 mL)	Urine (See Inst.)	APTIMA Urine Transport Tube	Room Temperature

Instructions Specimen: First-void urine in Aptima® urine transport

180051

Volume: 2 mL of a 20 to 30 mL urine collection

Container: Aptima® urine transport

Collection: Urine specimen: The patient should not have urinated for at least one hour prior to specimen collection. Direct patient to provide a first-catch urine (approximately 20 to 30 mL of the initial urine stream) into a urine collection cup free of any preservatives. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity; lesser volumes may not adequately rinse organisms into the specimen. Female patients should not cleanse the labial area prior to providing the specimen. Add urine to the Aptima® Combo 2 urine collection device. The final volume must be between the two black lines on the device (about 2 mL).

Storage Instructions: Maintain specimen at room temperature or refrigerate (2°C to 30°C). StabilityRoom temperature 30 days (stability provided by manufacturer or literature reference)

GENERAL INFORMATION

Expected TAT 3-4 days

Performing Labcorp Test

Code

Notes

RESULT CODE	RESULT NAME	LOINC
6907667	M genitalia Ur	88226-6
5195379	M. hominis Urine	68546-1
5195380	Ureaplasma Ur	68547-9
6910151	Chlamydia Ur	42931-6
6910155	Gonorrhea Ur	60256-5

CPT Code(s) 87491, 87591, 87798x2, 87563

Lab Section Reference Lab

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025