Copper, Plasma and RBC

Order Name: Copper, PI/RBC Test Number: 5195281 Revision Date: 09/30/2024

TEST NAME		МЕТ	HODOLOGY	LOINC CODE
Copper, Plasma			ntitative Inductively Coupled Plasma-Mass Sp	pectrometry 5631-7
Copper, RBC	Inductively-Coupled Plasma/Mass Spectrometry		5630-9	
SPECIMEN REQUIR	EMENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	3 mL (0.5 mL)	Packed RBCs and plasma	Royal Blue - EDTA (Trace Element Free)	Refrigerated
Instructions	Specimen: BOTH 3mL(0.5mL) Packed RBCs and 3mL(0.5mL) Plasma Keep Refrigerated. Container: Royal blue-top (EDTA) metal-free tube Collection: Both packed cells and plasma are required. Separate before shipping. Failure to separate may cause rejection due to hemolysis of red cells and contamination of the plasma. Transfer RBCs and plasma into separate plastic, acid-washed, or trace metal-free tubes. Specimen must be received at the test facility within five days of collection.			

Causes for Rejection: Failure to separate plasma and red cells; specimen received more than five days after collection

GENERAL INFORMATION		
Expected TAT	4 - 6 days	
Clinical Use	Evaluate copper stores and accumulation	
Performing Labcorp Test Code	081162	
CPT Code(s)	82525x2	
Lab Section	Reference Lab	