Coumadin (Warfarin) Level

Order Name: Coumadin
Test Number: 3606450
Revision Date: 08/31/2023

		METHODOLOGY	LOINC CODE
) Level		HPLC/Tandem Mass Spectrometry	4098-0
REMENTS			
Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
1 mL (0.3 mL)	Serum	Clot Activator (Red Top, No-Gel)	Room Temperature
1 mL (0.3 mL)	Plasma	EDTA (Lavender Top)	Room Temperature
Specimen Type: Red-top tube, lavender-top (EDTA) tube DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel. Specimen Collection: Promptly centrifuge and separate serum or plasma into a plastic screw capped vial using approved guidelines.			
	Specimen Volume (min) 1 mL (0.3 mL) 1 mL (0.3 mL) Specimen Type: Red-top tuber The use of gel-barrier tubes is Specimen Collection: Promp	Specimen Volume (min) Specimen Type 1 mL (0.3 mL) Serum 1 mL (0.3 mL) Plasma Specimen Type: Red-top tube,lavender-top (EDTA) tube DC The use of gel-barrier tubes is not recommended due to slow Specimen Collection: Promptly centrifuge and separate ser	Specimen Volume (min) Specimen Type Specimen Container 1 mL (0.3 mL) Serum Clot Activator (Red Top, No-Gel) 1 mL (0.3 mL) Plasma EDTA (Lavender Top) Specimen Type: Red-top tube,lavender-top (EDTA) tube DO NOT USE A GEL-BARRIER TUBE. The use of gel-barrier tubes is not recommended due to slow absorption of the drug by the gel.

GENERAL INFORMATION		
Expected TAT	3 - 4 days	
Performing Labcorp Test Code	832122	
CPT Code(s)	80375	
Lab Section	Reference Lab	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025