Reptilase Time with reflex to Reptilase Time 1:1 Mix

Order Name: **REPTILASE TIME**

Test Number: 6905511 Revision Date: 12/10/2022

TEST NAME METHODOLOGY LOINC CODE

Reptilase Time with reflex to Reptilase Time 1:1 Mix

Electromagnetic Mechanical Clot Detection

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL	Plasma	Sodium Citrate 3.2% (Blue Top)	Frozen
Instructions	Specimen Type: Blue-top (sodium citrate) tube Specimen Storage: FREEZE Specimen Collection: CITRATED PLASMA SAMPLES SHOULD BE COLLECTED BY DOUBLE CENTRIFUGATION. Blood should be collected in a blue-top tube containing 3.2% buffered sodium citrate.[1] Evacuated collection tubes must be filled to completion to ensure a proper blood to anticoagulant ratio.[2,3] The sample should be mixed immediately by gentle inversion at least six times to ensure adequate mixing of the anticoagulant with the blood. A discard tube is not required prior to collection of coagulation samples except when using a winged blood collection device (i.e. 'butterfly'), in which case a discard tube should be used.[4,5] When noncitrate tubes are collected for other tests, collect sterile and nonadditive (red-top) tubes prior to citrate (blue-top) tubes. Any tube containing an alternate anticoagulant should be collected after the blue- top tube. Gel-barrier tubes and serum tubes with clot initiators should also be collected after the citrate tubes. Centrifuge for 10 minutes and carefully remove 2/3 of the plasma using a plastic transfer pipette, being careful not to disturb the cells. Deliver to a plastic transport tube, cap, and recentifuge for 10 minutes. Use a second plastic pipette to remove plasma, staying clear of the platelets at the bottom of the tube. Transfer the plasma into a LabCorp PP transpak frozen purple tube with screw cap (LabCorp 49482). FREEZE immediately and maintain frozen until tested. Please print and use the Volume Guide for Coagulation Testing to ensure proper draw volume.			o ensure a proper blood to adequate mixing of the anting a winged blood collection device sts, collect sterile and nonadditive after the blue- top tube. Gel-barrier and carefully remove 2/3 of the and recentifuge for 10 minutes. The plasma into a LabCorp PP and Please print and use the

GENERAL INFORMATION		
Expected TAT	2 - 3 days	
Performing Labcorp Test Code	117180	
Notes	Labcorp Test Code: 117180	
CPT Code(s)	85635	
Lab Section	Reference Lab	

Specimen Stability: Ambient: Not Available, Refrigerated : Not Available, Frozen: Not Available

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025